



Received on 05 March 2020; received in revised form, 23 June 2020; accepted, 30 June 2020; published 01 March 2021

STUDIES ON GC-MS PROFILING OF SOME SEAWEEDS OF MAHIM BEACH DIST. PALGHAR, MAHARASHTRA USING VARIOUS SOLVENTS

Viraj Chabake ¹, Sakshi Chaubal ^{*2} and Tanmay Patil ¹

Department of Botany ¹, The Institute of Science, Mumbai - 400032, Maharashtra, India.

Department of Botany ², Mah. Govt's Elphinstone College Mumbai - 400032, Maharashtra, India.

Keywords:

Bioactive compounds, Crude extract, GC-MS analysis, Seaweeds etc.

Correspondence to Author:

Sakshi Chaubal

Associate Professor,
Department of Botany,
Mah. Govt's Elphinstone College
Mumbai - 400032, Maharashtra,
India.

E-mail: sakshi_chaubal@rediffmail.com

ABSTRACT: Objective: To investigate the bioactive constituents of *Ulva lactuca*, *Sargassum prismaticum*, and *Sarconema filiforme* using gas chromatography-mass spectrometry (GC-MS) analysis. **Methods:** 10 gm sample of each seaweed was successfully extracted with HPLC grade methanol and pet. ether with continuous shaking in a dark room for 36 h. The concentrated extracts were subjected to GC MS analysis. **Results:** The GC MS analysis of methanol and pet ether extracts of *U. lactuca*, *S. prismaticum* and *S. filiforme* show various types of phytochemicals screened under different R.T. Major compounds were 9-Octadecenoic acid (Z)-, phenyl-methyl ester, Stannane, 1,3-dithian-2-ylidenebis[tributyl], Octadecane, 3-ethyl-5-(2-ethylbutyl) in methanol extract and Decane, 6-ethyl-2-methyl-, Tricosane and Sulfurous acid, 2-ethylhexyl hexyl ester in pet. ether. These bioactive compounds are considered as biologically and pharmacologically important. **Conclusion:** The three different extracts contain diverse bioactive compounds that were identified and characterized spectroscopically. Hence, the identification of different biologically active compounds in the extracts of seaweeds justifies further biological and pharmacological studies.

INTRODUCTION: Seaweeds are one of the most significant existing marine resources in the world. These macroalgae have been a good source of food, fodder, manure, and medicine in the east as well as in the west, since centuries ^{1,2}. Seaweeds have been classified into three major divisions, namely Chlorophyta (Green algae), Phaeophyta (Brown algae), and Rhodophyta (red algae) based on their photosynthetic pigments and reserve food ³. Seaweeds are a significant marine resource as they are able to produce a great variety of secondary metabolites that could lead to the development of new functional ingredients ⁴.

Seaweeds comprise of large number of structurally diverse bioactive compounds such as alkaloids, glycosides, saponins, tannins, flavonoids, steroids etc., which are of immense medicinal value and use in a broad spectrum of biological activities ⁵.

All over the world, seaweeds have been studied and found to have enormous therapeutic potential as they possess anti-oxidant, neuroprotective, anti-fungal, anti-viral anti-inflammatory, anticancer, anti-HIV, antimutagenic, scavenging free radicals and antimicrobial activities. Seaweeds are also good source of vitamins such as A, B₁, B₁₂, C, D, E, riboflavin, niacin, pantothenic acid and folic acid as well as minerals such as Ca, P, Na, and K ⁶. Variation in nutrient content and bioactive compounds is observed in seaweeds in different seasonal cycles and with different environmental factors like ecological distribution, salinity, temperature, light, etc. ^{7, 8, 9, 10}

<p>QUICK RESPONSE CODE</p> 	<p>DOI: 10.13040/IJPSR.0975-8232.12(3).1644-50</p> <p>This article can be accessed online on www.ijpsr.com</p> <p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.12(3).1644-50</p>
-----------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

In spite of their significance towards mankind, seaweeds are relatively ignored in India. India has a total geographical area of about 329 million hectares with a vast coastline of over 7000 km^{11, 12}. Along the coast of India, many of the rocky beaches, coral reefs, lagoons, estuaries, and mudflats provide ideal habitats for the growth of marine algae¹³. The marine algae resources are abundant around coastlines of Mumbai, Palghar, Raigad, Ratnagiri, and Sindhudurg in Maharashtra¹⁴. The distribution of seaweed resources along the Indian coast was first mapped by Thivy¹⁵.

Large numbers of bioactive compounds have been isolated from the seaweeds so far; still, there is a need to explore and analyze many more seaweeds of different places which might show variation in biochemical content and could be used to produce various food supplements, industrial products, manure and compounds of pharmaceutical interest to cure different ailments.

Mahim beach, Dist. Palghar is a rocky beach. It shows ample collection of several seaweeds. Despite wealth of valuable raw products, seaweeds of this beach remained largely unexplored. *Ulva lactuca*, *Sargassum prismaticum*, and *Sarconema filiforme* were abundant at the Mahim beach and thus were selected for the present study of some bioactive compounds contained in them. *Ulva lactuca* belonging to the division Chlorophyta is a potential source of polysaccharides, proteins, pigments, cellulose, amino acids, lipids, fatty acids, and minerals, etc.^{16, 17} It has the capacity to rapidly proliferate and stated to exhibit a wide range of applications in food and health industry. Significant therapeutic properties like antibacterial, antifungal, antioxidant etc., are reported from *Ulva* sp. collected all over the world. *Sarconema filiforme* belonging to division Rhodophyta has considerable pharmacological, antimicrobial, antifungal, and cytotoxic potentials¹⁸.

Few sps. of *Sargassum* (Phaeophyta) are reported to be good source of phytochemical contents such as carotenoids, terpenes, phenolic compounds, fatty acids, lipids, and amino acids¹⁹. GC MS technique was selected for the characterization as it is the most appropriate technique to analyze and identify bioactive components of alcohols, esters, fatty acids, amino acids, steroids, long-chain hydro-

carbons, essential oils, etc. Crude seaweed extract was made using solvents of different polarities.

Studies related to bioactive compounds of seaweeds of Mahim beach, Dist Palghar are reported for the first time.

MATERIALS AND METHODS:

Collection of Seaweeds: Fresh seaweed *Ulva lactuca*, *Sargassum prismaticum*, and *Sarconema filiforme* were collected from intertidal regions of Mahim beach, dist. Palghar during the month of December 2017. The collected seaweed sample was cleaned with the seawater until unwanted impurities and adhering sand particles were removed. It was then cleaned thoroughly with tap water to remove the salty surface materials. The seaweed sample was shade dried for 7-8 days. The dried seaweed samples were powdered using mixer and passed through 0.5 mm sieve. It was stored in refrigerator for further study.

Seaweed Extraction: Seaweed extracts were obtained using solvents of different polarities. The seaweed powdered sample (10 gm) was extracted successively with HPLC grade methanol and pet ether using the cold extraction method. The sample was kept in a dark room for 36 hrs with continuous shaking. The extract was collected and filtered using Whatman no.1. It was evaporated to dryness by a rotavap. The final residue obtained was stored at 4 °C until further use. The volatile bioactive compounds present in methanol and pet. ether extracts of the seaweeds were identified by GC-MS characterization.

GC-MS Instruments: GC MS analysis of a different solvent extract of *Ulva lactuca*, *Sargassum prismaticum*, and *Sarconema filiforme* was carried out with Agilent 7890 system and gas chromatography interfaced to a mass spectrometer (GC-MS) employing the following conditions: HP 5 column [(5%-Phenyl)-methylpolysiloxane] (30mm × 0.25mm × 0.25µm] thickness, Helium gas (99.999%) was used as a carrier gas at a constant flow of 1ml/min, and injection volume of 1µl was employed (split ratio of 10:1) operating in electron impact mode at 70eV; injector temperature 250 °C; Ion-source temperature 280 °C. The oven temperature was programmed from 80 °C (isothermal for 1 min), with an increase of 10 °C

/min, to 200 °C then 5 °C/min to 280 °C ending with 9 min, isothermal at 280 °C. Mass spectra were taken at 70eV; a scan-interval of 0.5 seconds and fragments vary from 45-550 Da. Total GC running time was 35 min.

Identification of Chemical Constituents: The resulting peaks were analyzed with the database of the National Institute Standard and Technology (NIST) a library, which has more than 62,000 patterns. The spectrum of the unknown component was compared with the spectrum of the known components stored in it. The Name, Molecular weight and structure of the components of the seaweed sample was determined²⁰.

RESULTS: The volatile bioactive compounds present in methanol and pet. ether extract of *Ulva lactuca*, *Sargassum prismaticum*, and *Sarconema filiforme* were identified by GC-MS analysis. The methanol and pet. ether extracts of seaweeds indicated the presence of different types of phytocompounds **Table 1-2** screened under different retention times (R.T). A total of (9, 4 and



9 of methanol extract) and (4, 7, and 11 of pet. ether extract) were observed with retention times for *U. lactuca*, *S. prismaticum* and *S. filiforme* respectively, as shown in **Fig. 1-6**. Chemical constituents were identified using spectral database NIST 11 software installed in the GC-MS. The compounds prediction is based on Pubchem²⁰. The compound name with R.T, molecular formula, molecular weight and concentration % in the methanol and petether extract of seaweeds *Ulva lactuca*, *Sargassum prismaticum* and *Sarconema filiforme* are presented in **Table 1-2**. 9-Octadecenoic acid (Z)-, phenylmethyl ester are most abundant (R.T 31.81, 31.52%) in methanol *U. lactuca* extract followed by Stannane, 1,3-dithian-2-ylidenebis[tributyl- (R.T 14.82, 70.59%) in *S. prismaticum* and Octadecane, 3-ethyl-5-(2-ethylbutyl)- (R.T 31.62, 36.50%) present in *S. filiforme*. Crude extract of petether indicates most abundant compound as Decane, 6-ethyl-2-methyl- (R.T 31.86, 41.07%), Tricosane (R.T 31.66, 25.84 %) and Sulfurous acid, 2-ethylhexyl hexyl ester (R.T 19.42, 31.96%) respectively.

TABLE 1: COMPOUNDS IDENTIFIED IN THE METHANOL EXTRACT OF *ULVA LACTUCA*, *SARGASSUM PRISMATICUM* AND *SARCONEMA FILIFORME* BY GC-MS ANALYSIS

Extract	RT (min)	Name of compound	Molecular formula	Molecular weight (g/mol)	Peak %
<i>Ulva</i> methanol	15.65	Decanoic acid 2-methyl-	C ₁₁ H ₂₂ O ₂	186	3.288906
	18.93	11,14,17-eicosatrienoic acid,methyl ester	C ₂₁ H ₃₆ O ₂	320	8.680821
	22.67	1-Nonyne,-/methyl	C ₁₀ H ₁₈	138	1.552746
	22.97	9-Oxabicyclo[4.3.0]non-2-ene-4,5-dicarboxylic,8-hydroxy-1,2,7,7-tetramethyl-,dimethyl ester	C ₁₆ H ₂₄ O ₆	312	14.59502
	23.25	Silane, (1,3,5-benzenetriyltris(methylene)tris(trimethyl-2,2'-Thiobis[4-methyl-6-(1, 1, 3, 3-tetramethylbutyl)phenol]	C ₁₅ H ₃₆ Si ₃	294	18.11925
	23.49	(t-butyl-dimethylsilyl)[2-methyl-2-(4-methyl-pent-3-enyl)-cyclopropyl]-methanol	C ₃₀ H ₄₆ O ₂	470	9.687301
	23.86	d-Mannitol, 1-decylsulfonyl-	C ₁₇ H ₃₄ OSi	282.537	1.73179
	31.26	9-Octadecenoic acid(Z)-, phenylmethyl ester	C ₁₆ H ₃₄ O ₇ S	370.501	10.82201
	31.81	1,14-Tetradecanediol	C ₂₅ H ₄₀ O ₂	372	31.52216
<i>Sargassum</i> methanol	14.15	Stannane,1,3-dithian-2-ylidenebis[tributyl	C ₁₄ H ₃₀ O ₂	230.392	5.847037
	14.82	Probuocol	C ₂₈ H ₆₀ S ₂ S _{n2}	700	70.59108
	29.32	Tetradecane, 1-iodo-	C ₃₁ H ₄₈ O ₂ S ₂	516.843	5.945797
<i>Sarconema</i> methanol	31.85	1-Nitrododecane	C ₁₄ H ₂₉ I	324.2845	17.61608
	22.75	Eseroline,octylcarbamate(ester)	C ₁₂ H ₂₅ NO ₂	215.337	0.034011947
	23.07	1,2-Benzenedicarboxylic acid, ditridecyl ester;	C ₂₂ H ₃₅ N ₃ O ₂	373	0.654931203
	23.39	Sulfurous acid, 2-propyl tridecyl ester	C ₃₄ H ₅₈ O ₄	530.8219	1.159361947
	23.74	1-Bromoeicosane	C ₁₆ H ₃₄ O ₃ S	306	1.281223925
	24.78	Octadecane, 3-ethyl-5-(2-ethylbutyl)-	C ₂₀ H ₄₁ Br	361.452	6.102793402
	26.90	Octadecane, 3-ethyl-5-(2-ethylbutyl)-	C ₂₆ H ₅₄	366	19.3288882
	27.95	Cholest-5-en-3-ol, 24-propylidene-, (3 β)-	C ₂₆ H ₅₄	366.7070	13.65299758
28.7	Octadecane, 3-ethyl-5-(2-ethylbutyl)-	C ₃₀ H ₅₀ O	426	21.27953871	
31.62	Octadecane, 3-ethyl-5-(2-ethylbutyl)-	C ₂₆ H ₅₄	366.7070	36.50625308	

TABLE 2: COMPOUNDS IDENTIFIED IN THE PET. ETHER EXTRACT OF *ULVA LACTUCA*, *SARGASSUM PRISMATICUM* AND *SARCONEMA FILIFORME* BY GC-MS ANALYSIS

Extract	RT (min)	Name of compound	Molecular formula	Molecular weight (g/mol)	Peak %
<i>Ulva</i> Pet. ether	19.44	Pentanoic acid, 5-hydroxy-, 2,4-di-t-butylphenyl esters	C ₁₉ H ₃₀ O ₃	306.446	36.37524
	19.99	Tetradecane, 1-iodo-	C ₁₄ H ₂₉ I	324.29	1.34168
	31.40	L-Norvaline, N-(2-methoxyethoxycarbonyl)-, isoheptyl ester	C ₁₅ H ₂₉ NO ₅	303	21.20727
<i>Sargassum</i> Pet. ether	31.86	Decane, 6-ethyl-2-methyl-	C ₁₃ H ₂₈	184	41.07581
	9.94	Sulfurous acid, 2-ethylhexyl hexyl ester	C ₁₄ H ₃₀ O ₃ S	278.451	12.60196
	12.35	1-Hentetracontanol	C ₄₁ H ₈₄ O	593.122	4.817065
	19.33	L-Norvaline, N-(2-methoxyethoxycarbonyl)-, undecyl ester	C ₂₀ H ₃₉ NO ₅	373.52736	0.580089
	19.45	L-Norvaline, N-(2-methoxyethoxycarbonyl)-, undecyl ester	C ₂₀ H ₃₉ NO ₅	373.52736	29.55033
	19.97	6-Fluoro-2-trifluoromethylbenzoic acid, 2-formyl-4,6-dichlorophenyl ester	C ₁₅ H ₆ Cl ₂ F ₄ O ₃	381.106	1.319213
<i>Sarconema</i> Pet. ether	31.66	Tricosane	C ₂₃ H ₄₈	324.637	25.84063
	31.86	1-Octadecanesulfonyl chloride	C ₁₈ H ₃₇ ClO ₂ S	353.002	25.29071
	9.3	Decane, 2,9-dimethyl-	C ₁₂ H ₂₆	170.3348	6.073613
	12.03	6-Fluoro-2-trifluoromethylbenzoic acid, 2-formyl-4,6-dichlorophenyl ester	C ₁₅ H ₆ Cl ₂ F ₄ O ₃	381.106	0.478575
	12.25	Sulfurous acid, 2-ethylhexyl hexyl ester	C ₁₄ H ₃₀ O ₃ S	278.451	20.24306
	19.18	Decane, 1-bromo-2-methyl-	C ₁₁ H ₂₃ Br	235.209	1.647081
	19.42	Sulfurous acid, 2-ethylhexyl isoheptyl ester	C ₁₄ H ₃₀ O ₃ S	278.451	31.96578
	19.79	l-norvaline, n-(2-methoxyethoxycarbonyl)-, octyl ester	C ₁₇ H ₃₃ NO ₅	331	1.844492
	19.97	l-Valine, n-pentafluoropropionyl-, undecyl ester	C ₁₉ H ₃₂ F ₅ NO ₃	417.4543	1.48032
	27.23	Octadecane, 3-ethyl-5-(2-ethylbutyl)-	C ₂₆ H ₅₄	366.7070	3.714758
	28.51	Di-n-decylsulfone	C ₂₀ H ₄₂ O ₂ S	346.614	4.630403
31.54	Heptacosane	C ₂₇ H ₅₆	380.745	8.254636	
31.85	Octadecane, 3-ethyl-5-(2-ethylbutyl)-	C ₂₆ H ₅₄	366.7070	19.66728	

**FIG. 1: GC-MS CHROMATOGRAM OF THE CRUDE METHANOL EXTRACT OF *ULVA LACTUCA*****FIG. 2: GC-MS CHROMATOGRAM OF THE CRUDE METHANOL EXTRACT OF *SARGASSUM PRISMATICUM***




FIG. 3: GC-MS CHROMATOGRAM OF THE CRUDE METHANOL EXTRACT OF *SARCONEMA FILIFORME*




FIG. 4: GC-MS CHROMATOGRAM OF THE CRUDE PET. ETHER EXTRACT OF *ULVA LACTUCA*




FIG. 5: GC-MS CHROMATOGRAM OF THE CRUDE PET. ETHER EXTRACT OF *SARGASSUM PRISMATICUM*




FIG. 6: GC-MS CHROMATOGRAM OF THE CRUDE PET. ETHER EXTRACT OF *SARCONEMA FILIFORME*

DISCUSSION: Marine algae have been rich source of carotenoids, polyunsaturated fatty acids²¹, polysaccharides²², flavonoids, phenolic acids²³, sterols²⁴ Phlorotannins²⁵ etc. Present GC-MS analysis of crude extracts of methanol and pet. ether of *Ulva lactuca*, *Sargassum prismaticum* and *Sarconema filiforme* revealed various compounds. Major compounds were 9-Octadecenoic acid(Z)-, phenylmethyl ester, Stannane, 1,3-dithian-2-ylidenebis [tributyl, Octadecane, 3-ethyl-5-(2-ethylbutyl) in methanol extract and Decane, 6-ethyl-2-methyl-, Tricosane and Sulfurous acid, 2-ethylhexyl hexyl ester in pet. ether. Some other important bioactive compounds mentioned in **Table 1-2** are rich source of structurally novel and biologically active metabolites and can be used as potential natural antioxidants in different food and pharmaceutical products²⁶. Babu et al., (2014) reported 17 different components in ethanolic extract of *Ulva lactuca*. Major components Phytol (10.51%), Hexadecanoic acid, ethyl ester (20.24%) and (E)-9- Octadecenoic acid ethyl ester (9.59%) are considered to have significant medicinal properties. Kumar et al., (2013) found 12 compounds having antibacterial activity in methanolic extract of *Sargassum tenerrimum* by GC MS analysis. Red algae *Halimeda opuntia* and *Sarconema filiforme* are described as potent algal species with considerable antimicrobial, antiplasmid, and cytotoxic activity²⁹.

CONCLUSION: The present study helps to predict the formula and structure of bioactive compounds in *Ulva lactuca*, *Sargassum prismaticum*, and *Sarconema filiforme*. Further work related to verifying their efficacy may lead to the improvement of drug formulation.

ACKNOWLEDGEMENT: Nil

CONFLICTS OF INTEREST: Nil

REFERENCES:

- Chapman VJ and Chapman DJ: Seaweeds and their uses, Chapman and Hall, New York, Edn 3 1980; 334.
- Arasaki S and Arasaki T: Vegetables from the Sea. Japan Pub. Inc, Tokyo 1983; 196.
- Dawczynski C, Schubert R and Jahreis G: Amino acids, fatty acids, and dietary fibre in edible seaweed products. Food Chemistry 2007; 103: 891-99.
- Freile-Pelegrin Y, Robledo D, Chan-Bacab MJ and Ortega-Morales BO: Antileishmanial properties of tropical marine algae extracts. Fitoterapia 2008; 79(5): 374-77.
- Maria JP, Falque E and Dominguez H: Antimicrobial action of compounds from marine seaweed. Mar Drugs 2016; 14(3): 52.
- Dhargalkar VK and Pereira N: Seaweed: promising plant of the millennium. Science and Culture 2005; 71: 60-66.
- Kailas AP and Nair MS: Saponins and the *in-vitro* bioactivities of different solvent extracts of some tropical green and red seaweeds. J Coast Lif Med 2015; 3: 931-43.
- Gaubert J, Payri CE, Vieira C, Solanki H and Thomas OP: High metabolic variation for seaweeds in response to environmental changes: A case study of the brown algae *Lobophora* in coral reefs. Sci Rep 2019; 9: 1-12.
- Dhevi S, Deivasigamani B. Phytochemical profiling and GC-MS analysis of *Caulerpa racemosa*. Res J Life Sci Bioinform Pharm Chem Sci 2018; 4: 155-65.
- Kavitha J and Palani S: Phytochemical screening, GC-MS analysis and antioxidant activity of marine algae *Chlorococcum humicola*. World Journal of Pharmacy and Pharmaceutical Sciences 2016; 5(6): 1154-67.
- Anonymous: Status of plant diversity in india.org [ENVIS Centre on Floral Diversity]. India: Central National Herbarium, Botanical Survey of India, Howrah. Available from: http://www.bsienviis.nic.in/Database/Status_of_Plant_Diversity_in_India_17566.aspx updated 12/10/2017.
- Sahoo D, Sahu N and Sahoo A: Critical survey of seaweed diversity on Chilika Lake, India. Algae 2003; 18: 1-12.
- Rajkumar G and Bhavan PS: Phytochemical characterization of the marine brown alga *Turbinaria ornata*. Res J Chem Environ 2017; 21: 54-63.
- Deodhar HD: The biology of marine algae of Bombay, Ph.D thesis, Savitribai Phule Pune University, Pune, India 1989.
- Thivy F: Economic seaweeds. In Fisheries of West Coast of India, Bangalore, Current Science 1958; 74-80.
- Kumar JGS, Umamaheswari S, Kavimani S and Ilavarasan R: Pharmacological potential of green algae *Caulerpa*: A review. Int J Pharm Sci Res 2019; 10: 1014-24.
- Ortiz J, Romero N, Robert P, Araya J, Lopez-Hernandez J, Bozzo C, Navarete E, Osorio A and Rios A: Dietary fiber, amino acid, fatty acid and tocopherol contents of the edible seaweeds *Ulva lactuca* and *Durvillaea antarctica*. Food Chem 2006; 99: 98-104.
- Samy A and Selim S: Antimicrobial, antiplasmid and cytotoxicity potentials of marine algae *Halimeda opuntia* and *Sarconema filiforme* collected from Red Sea Coast. WASET 2012; 1: 61.
- Hossain Z, Kurihara H and Takahashi K: Biochemical composition and lipid compositional properties of the brown alga *Sargassum horneri*. Pakistan Journal of Biological Sciences 2003; 6: 1497-1500.
- Anonymous: National Institutes of Health (NIH) [Pubchem] U.S National Library of medicine Available from: <https://pubchem.ncbi.nlm.nih.gov/>.
- Pereira H, Barreira L, Figueiredo F, Custodio L, Vizetto Duarte C, Polo C, Resek E, Engelen A and Varela J: Polyunsaturated fatty acids of marine macroalgae: potential for nutritional and pharmaceutical applications. Mar Drugs 2012; 10: 1920-35.
- Raposo MFJ, Morais AMB and Morais RMS: Marine Polysaccharides from Algae with Potential Biomedical Applications. Mar Drugs 2015; 13: 2967-3028.
- Tibbetts SM, Milley JE and Lall SP: Nutritional quality of some wild and cultivated seaweeds: nutrient composition, total phenolic content and *in-vitro* digestibility. J Appl Phycol 2016; 26: 3575-85.
- Wells ML, Potin P, Craigie JS, Raven JA, Merchant SS, Helliwell KE, Smith AG, Camire ME and Brawley SH:

- Algae as nutritional and functional food sources: revisiting our understanding. *J Appl Phycol* 2017; 29: 949-82.
25. Thomas NV and Kim SK: Potential pharmacological applications of polyphenolic derivatives from marine brown algae. *Environ. Toxicol Pharmacol* 2011; 32: 325-35.
26. MMEI-Din S and MDEI-Ah A: Bioactivity and phytochemical constituents of marine red seaweeds (*Jania rubens*, *Corallina mediterranea* and *Pterocladia capillacea*). *Journal of Taibah University for Science* 2016; 10: 471-84.
27. Babu MJ, Raja DP, raj Arockia AA and Vinnarasi J: Chemical constituents and their biological activity of *Ulva lactuca* Linn. *International Journal of Pharmaceutical and Drug Analysis* 2014; 2(7): 595-600.
28. Kumar P, Senthamilselvi S and Govindaraju M: GC-MS profiling and antibacterial activity of *Sargassum tenerrimum*. *J of Pharmacy Research* 2013; 6: 88-92.
29. Selim SA: Antimicrobial, antiplasmid and cytotoxicity potentials of marine algae *Halimeda opuntia* and *Sarconema filiforme* collected from Red Sea Coast. *World Acad Sci Eng Technol* 2012; 61: 1154-9.

How to cite this article:

Chabake V, Chaubal S and Patil T: Studies on gc-ms profiling of some seaweeds of Mahim Beach Dist. Palghar, Maharashtra using various solvents. *Int J Pharm Sci & Res* 2021; 12(3): 1644-50. doi: 10.13040/IJPSR.0975-8232.12(3).1644-50.

All © 2013 are reserved by the International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)